

A PRESENTATION OF THE DEFORMED $W_{1+\infty}$ ALGEBRA

N. ARBESFELD AND O. SCHIFFMANN

ABSTRACT. We provide a generators and relation description of the deformed $W_{1+\infty}$ -algebra introduced in previous joint work of E. Vasserot and the second author. This gives a presentation of the (spherical) cohomological Hall algebra of the one-loop quiver, or alternatively of the spherical degenerate double affine Hecke algebra of $GL(\infty)$.

INTRODUCTION

In the course of their work on the cohomology of the moduli space of $U(r)$ -instantons on $\mathbb{P}^2$ in relation to W -algebras and the AGT conjecture (see [SV]) E. Vasserot and the second author introduced a certain one-parameter deformation $\mathbf{SH}^c$ of the enveloping algebra of the Lie algebra $W_{1+\infty}$ of algebraic differential operators on $\mathbb{C}^*$. The algebra $\mathbf{SH}^c$ —which is defined in terms of Cherednik’s double affine Hecke algebras—acts on the above mentioned cohomology spaces (with a central character depending on the rank n of the instanton space). For the same value of the central character, $\mathbf{SH}^c$ is also strongly related to the affine W algebra of type $\mathfrak{gl}_n$, and has the same representation theory (of admissible modules) as the latter. The same algebra $\mathbf{SH}^c$ arises again as the (spherical) cohomological Hall algebra of the quiver with one vertex and one loop, and as a degeneration of the (spherical) elliptic Hall algebra (see [SV, Sec. 4, 8]. It also independently appears in the work of Maulik and Okounkov on the AGT conjecture, see [MO].

The definition of $\mathbf{SH}^c$ given in [SV] is in terms of a stable limit of spherical degenerate double affine Hecke algebras, and does not yield a presentation by generators and relations. In this note, we provide such a presentation, which bears some resemblance with Drinfeld’s new realization of quantum affine algebras and Yangians. Namely, we show that $\mathbf{SH}^c$ is generated by families of elements in degrees $-1, 0, 1$, modulo some simple quadratic and cubic relations (see Theorems 3.1, 3.2).

The definition of $\mathbf{SH}^c$ is recalled in Section 1. In the short Section 2 we briefly recall the links between $\mathbf{SH}^c$ and Cherednik algebras, resp. W -algebras. The presentation of $\mathbf{SH}^c$ is given in Section 3, and proved in Section 4. Although we have tried to make this note as self-contained as possible, there are multiple references to statements in [SV] and the reader is advised to consult that paper (especially Sections 1 and 8) for details.

1. DEFINITION OF $\mathbf{SH}^c$

1.1. Symmetric functions and Sekiguchi operators. Let κ be a formal parameter, and let us set $F = \mathbb{C}(\kappa)$. Let us denote by Λ_F the ring of symmetric polynomials in infinitely many variables with coefficients in F , i.e.

$$\Lambda_F = F[X_1, X_2, \dots]^{\mathfrak{S}_\infty} = F[p_1, p_2, \dots].$$

For λ a partition, we denote by J_λ the integral form of the Jack polynomial associated to λ and to the parameter $\alpha = 1/\kappa$. It is well-known that $\{J_\lambda\}$ forms a basis of Λ_F (see e.g. [S], or [SV, Sec. 1.3, 1.6]).

The polynomials J_λ arise as the joint spectrum of a family of commuting differential operators $\{D_{0,l}\}, l \geq 1$ called Sekiguchi operators. In the above normalization, these may be characterized through the following relations :

$$(1.1) \quad D_{0,l}(J_\lambda) = \sum_{s \in \lambda} c(s)^{l-1} J_\lambda$$

where s runs through the set of boxes in the partition λ , and where $c(s) = x(s) - \kappa y(s)$ is the content of s . Here $x(s), y(s)$ denote the x and y -coordinates of the box s , when λ is drawn according to the continental convention (see [SV, Sec. 0.1]).

We denote by $D_{l,0} \in \text{End}(\Lambda_F)$ the operator of multiplication by the power-sum function p_l .

1.2. The algebras $\mathbf{SH}^+$ and $\mathbf{SH}^>$. Let $\mathbf{SH}^+$ be the unital subalgebra of $\text{End}(\Lambda_F)$ generated by $\{D_{0,l}, D_{l,0} \mid l \geq 1\}$. For $l \geq 1$ we set $D_{1,l} = [D_{0,l+1}, D_{1,0}]$. This relation is still valid when $l = 0$, and we furthermore have

$$(1.2) \quad [D_{0,l}, D_{1,k}] = D_{1,k+l-1} \quad l \geq 1, k \geq 0.$$

We denote by $\mathbf{SH}^>$ the unital subalgebra of $\mathbf{SH}^+$ generated by $\{D_{1,l} \mid l \geq 0\}$, and by $\mathbf{SH}^0$ the unital subalgebra of $\mathbf{SH}^+$ generated by the Sekiguchi operators $\{D_{0,l} \mid l \geq 1\}$. It is known (and easy to check from (1.1)) that the $D_{0,l}$ are algebraically independent, i.e. $\mathbf{SH}^0 = F[D_{0,1}, D_{0,2}, \dots]$.

Observe that by (1.2), the operators $\text{ad}(D_{0,l})$ preserve the subalgebra $\mathbf{SH}^>$. This allows us to view $\mathbf{SH}^+$ as a semi-direct product of $\mathbf{SH}^0$ and $\mathbf{SH}^>$. In fact, the multiplication map induces an isomorphism

$$(1.3) \quad \mathbf{SH}^> \otimes \mathbf{SH}^0 \simeq \mathbf{SH}^+$$

(see [SV, Prop. 1.18]).

1.3. Grading and filtration. The algebra $\mathbf{SH}^+$ carries an $\mathbb{N}$ -grading, defined by setting $D_{0,l}, D_{1,k}$ in degrees zero and one respectively. This grading, which corresponds to the degrees as operators on polynomials will be called the *rank* grading. It also carries an $\mathbb{N}$ -filtration compatible with the rank grading, induced from the filtration by the order of differential operators. It may be characterized as follows, see [SV, Prop. 1.2] : $\mathbf{SH}^+[\leq d]$ is the space of elements $u \in \mathbf{SH}^>$ satisfying

$$\text{ad}(z_1) \circ \dots \circ \text{ad}(z_{d+1})(u) = 0$$

for all $z_1, \dots, z_{d+1} \in F[D_{1,0}, D_{2,0}, \dots]$. We have $\mathbf{SH}^+[\leq 0] = F[D_{1,0}, D_{2,0}, \dots]$. The following is proved in [SV, Lemma 1.21]. Set $D_{r,d} = [D_{0,d+1}, D_{r,0}]$ for $r \geq 1, d \geq 0$.

Proposition 1.1. (i) *The associated graded algebra $\text{gr}\mathbf{SH}^+$ is equal to the free commutative polynomial algebra in the generators $D_{r,d} \in \text{gr}\mathbf{SH}^+[r, d]$, for $r \geq 0, d \geq 0, (r, d) \neq (0, 0)$.*

(ii) *The associated graded algebra $\text{gr}\mathbf{SH}^>$ is equal to the free commutative polynomial algebra in the generators $D_{r,d} \in \text{gr}\mathbf{SH}^+[r, d]$, for $r \geq 1, d \geq 0$.*

We will need the following slight variant of the above result, which can easily be deduced from [SV, Prop. 1.38]. For $r \geq 1$, set $D'_{r,d} = \text{ad}(D_{0,2})^d(D_{r,0})$. Then

$$(1.4) \quad D'_{r,d} \in r^{d-1} D_{r,d} \oplus \mathbf{SH}^>[r, \leq d-1].$$

In particular, $\text{gr}\mathbf{SH}^>$ is also freely generated by the elements $D'_{r,d} \in \text{gr}\mathbf{SH}^>[r, d]$.

1.4. The algebra $\mathbf{SH}^c$. Let $\mathbf{SH}^<$ be the opposite algebra of $\mathbf{SH}^>$. We denote the generator of $\mathbf{SH}^>$ corresponding to $D_{1,l}$ by $D_{-1,l}$. The algebra $\mathbf{SH}^c$ is generated by $\mathbf{SH}^>, \mathbf{SH}^0, \mathbf{SH}^<$ together with a family of central elements $\mathbf{c} = (c_0, c_1, \dots)$ indexed by $\mathbb{N}$, modulo a certain set of relations involving the commutators $[D_{-1,k}, D_{1,l}]$ (see [SV, Sec. 1. 8]). In order to write down these relations, we need a few notations. Set $\xi = 1 - \kappa$ and

$$\begin{aligned} G_0(s) &= -\log(s), & G_l(s) &= (s^{-1} - 1)/l, & l \geq 1, \\ \varphi_l(s) &= \sum_{q=1, -\xi, -\kappa} s^l (G_l(1 - qs) - G_l(1 + qs)), & l \geq 1, \end{aligned}$$

$$\phi_l(s) = s^l G_l(1 + \xi s).$$

We may now define $\mathbf{SH}^c$ as the algebra generated by $\mathbf{SH}^>, \mathbf{SH}^<, \mathbf{SH}^0$ and $F[c_0, c_1, \dots]$ modulo the following relations :

$$(1.5) \quad [D_{0,l}, D_{1,k}] = D_{1,k+l-1}, \quad [D_{-1,k}, D_{0,l}] = D_{-1,k+l-1},$$

$$(1.6) \quad [D_{-1,k}, D_{1,l}] = E_{k+l}, \quad l, k \geq 0,$$

where the elements E_h are determined through the formulas

$$(1.7) \quad 1 + \xi \sum_{l \geq 0} E_l s^{l+1} = \exp \left(\sum_{l \geq 0} (-1)^{l+1} c_l \phi_l(s) \right) \exp \left(\sum_{l \geq 0} D_{0,l+1} \varphi_l(s) \right).$$

Set $\mathbf{SH}^{0,c} = \mathbf{SH}^0 \otimes F[c_0, c_1, \dots]$. One can show that the multiplication map provides an isomorphism of F -vector spaces

$$\mathbf{SH}^> \otimes \mathbf{SH}^{0,c} \otimes \mathbf{SH}^< \simeq \mathbf{SH}^c.$$

Putting the generators $D_{\pm 1,k}$ in degree ± 1 and the generators $D_{0,l}, c_i$ in degree zero induces an $\mathbb{Z}$ -grading on $\mathbf{SH}^c$. One can show that the order filtration on $\mathbf{SH}^>, \mathbf{SH}^<$ can be extended to a filtration on the whole $\mathbf{SH}^c$, but we won't need this last fact.

2. LINK TO W-ALGEBRAS, CHEREDNIK ALGEBRAS AND SHUFFLE ALGEBRAS

2.1. Relation the Cherednik algebras. Let ω be a new formal parameter and let $\mathbf{SH}^\omega$ be the specialization of $\mathbf{SH}$ at $c_0 = 0, c_i = -\kappa^i \omega^i$. Let $\mathbf{H}_n$ be Cherednik's degenerate (or trigonometric) double affine Hecke algebra with parameter κ (see [C]). Let $\mathbf{SH}_n \subset \mathbf{H}_n$ be its spherical subalgebra. The following result shows that $\mathbf{SH}^\omega$ may be thought of as the stable limit of $\mathbf{SH}_n$ as n goes to infinity (see [SV, Sec. 1.7]) :

Theorem. *For any n there exists a surjective algebra homomorphism $\Phi_n : \mathbf{SH}^\omega \rightarrow \mathbf{SH}_n$ such that $\Phi_n(\omega) = n$. Moreover $\bigcap_n \text{Ker } \Phi_n = \{0\}$.*

2.2. Realization as a shuffle algebra. Consider the rational function

$$g(z) = \frac{h(z)}{z}, \quad h(z) = (z+1-\kappa)(z-1)(z+\kappa).$$

Following [FO], we may associate to $g(z)$ an $\mathbb{N}$ -graded associative F -algebra $A_{g(z)}$, the *symmetric shuffle algebra of $g(z)$* as follows. As a vector space,

$$A_{g(z)} = \bigoplus_{n \geq 0} F[z_1, \dots, z_n]^{\mathfrak{S}_n}$$

with multiplication given by

$$P(z_1, \dots, z_r) \star Q(z_1, \dots, z_s) = \sum_{\sigma \in Sh_{r,s}} \sigma \cdot \left(\prod_{\substack{1 \leq i \leq r \\ r+1 \leq j \leq r+s}} g(z_i - z_j) \cdot P(z_1, \dots, z_r) Q(z_{r+1}, \dots, z_{r+s}) \right)$$

where $Sh_{r,s} \subset \mathfrak{S}_{r+s}$ is the set of (r, s) shuffles inside the symmetric group $\mathfrak{S}_{r+s}$. Let $S_{g(z)} \subseteq A_{g(z)}$ denote the subalgebra generated by $A_{g(z)}[1] = F[z]$. The following is proved in [SV, Cor. 6.4] :

Theorem. *The assignment $S_{g(z)}[1] \ni z^l \mapsto D_{1,l}, l \geq 0$ induces an isomorphism of F -algebras*

$$S_{g(z)} \xrightarrow{\sim} \mathbf{SH}^>.$$

2.3. Relation to W -algebras. Let $W_{1+\infty}$ be the universal central extension of the Lie algebra of all differential operators on $\mathbb{C}^*$ (see e.g. [FKRW]). This is a $\mathbb{Z}$ -graded and $\mathbb{N}$ -filtered Lie algebra. The following result shows that $\mathbf{SH}$ may be thought of as a deformation of the universal enveloping algebra $U(W_{1+\infty})$ of $W_{1+\infty}$ (see [SV, App. F]) :

Theorem. *The specialization of $\mathbf{SH}^c$ at $\kappa = 1$ and $c_i = 0$ for $i \geq 1$ is isomorphic to $U(W_{1+\infty})$.*

More interesting is the fact that, for certain good choices of the parameters $c_0, c_1, \dots$, a suitable completion of $\mathbf{SH}^c$ is isomorphic to the current algebra of the (affine) W -algebra $W(\mathfrak{gl}_r)$ (see e.g. [A, Sec. 3.11]). Fix an integer $r \geq 1$, $k \in \mathbb{C}$ and let $(\varepsilon_1, \dots, \varepsilon_r)$ be new formal parameters. Let $\mathfrak{U}(W_k(\mathfrak{gl}_r))'$ be the formal current algebra of $W(\mathfrak{gl}_r)$ at level k , defined over the field $F(\varepsilon_1, \dots, \varepsilon_r)$ (see [SV, Sec. 8.4] for details). Let $\mathbf{SH}^{(r)}$ be the specialization of $\mathbf{SH}^c$ to $\kappa = k + r$, $c_i = \varepsilon_1^i + \dots + \varepsilon_r^i$ for $i \geq 0$. The following is proved in [SV, Cor. 8.24], to which we refer for details.

Theorem. *There is an embedding $\mathbf{SH}^{(r)} \rightarrow \mathfrak{U}(W_k(\mathfrak{gl}_r))'$ with a dense image, which induces an equivalence between the category of admissible $\mathbf{SH}^{(r)}$ -modules and the category of admissible $\mathfrak{U}(W_k(\mathfrak{gl}_r))'$ -modules.*

3. PRESENTATION OF $\mathbf{SH}^+$ AND $\mathbf{SH}^c$

3.1. Generators and relations for $\mathbf{SH}^+$. Consider the F -algebra $\widetilde{\mathbf{SH}}^+$ generated by elements $\{\tilde{D}_{0,l} \mid l \geq 1\}$ and $\{\tilde{D}_{1,k} \mid k \geq 0\}$ subject to the following set of relations :

$$(3.1) \quad [\tilde{D}_{0,l}, \tilde{D}_{0,k}] = 0, \quad \forall l, k \geq 1,$$

$$(3.2) \quad [\tilde{D}_{0,l}, \tilde{D}_{1,k}] = \tilde{D}_{1,l+k-1}, \quad \forall l \geq 1, k \geq 0,$$

$$(3.3) \quad (3[\tilde{D}_{1,2}, \tilde{D}_{1,1}] - [\tilde{D}_{1,3}, \tilde{D}_{1,0}] + [\tilde{D}_{1,1}, \tilde{D}_{1,0}]) + \kappa(\kappa - 1)(\tilde{D}_{1,0}^2 + [\tilde{D}_{1,1}, \tilde{D}_{1,0}]) = 0$$

$$(3.4) \quad [\tilde{D}_{1,0}, [\tilde{D}_{1,0}, \tilde{D}_{1,1}]] = 0.$$

Let $\widetilde{\mathbf{SH}}^0 = F[\tilde{D}_{0,1}, \tilde{D}_{0,2}, \dots]$ denote the subalgebra of $\widetilde{\mathbf{SH}}^+$ generated by $\tilde{D}_{0,l}, l \geq 1$, and let $\widetilde{\mathbf{SH}}^>$ be the subalgebra generated by $\tilde{D}_{1,k}, k \geq 0$. The algebras $\widetilde{\mathbf{SH}}^+, \widetilde{\mathbf{SH}}^0, \widetilde{\mathbf{SH}}^>$ are all $\mathbb{N}$ -graded, where $\tilde{D}_{0,l}$ and $\tilde{D}_{1,k}$ are placed in degrees zero and one respectively. According to the terminology used for $\mathbf{SH}^+$, we call this grading the *rank grading*.

Theorem 3.1. *The assignment $\tilde{D}_{0,l} \mapsto D_{0,l}, \tilde{D}_{1,k} \mapsto D_{1,k}$ for $l \geq 1, k \geq 0$ induces an isomorphism of graded F -algebras*

$$\phi : \widetilde{\mathbf{SH}}^+ \xrightarrow{\sim} \mathbf{SH}^+.$$

Obviously, the map ϕ restricts to isomorphisms $\widetilde{\mathbf{SH}}^0 \simeq \mathbf{SH}^0, \widetilde{\mathbf{SH}}^> \simeq \mathbf{SH}^>$. Note however that $\widetilde{\mathbf{SH}}^>$ is *not* generated by the elements $\tilde{D}_{1,k}$ with the sole relations (3.3, 3.4). Theorem 3.1 is proved in Section 4.

3.2. Generators and relations for $\mathbf{SH}^c$. For the reader's convenience, we write down the presentation of $\mathbf{SH}^c$, an immediate corollary of Theorem 3.1 above. Let $\widetilde{\mathbf{SH}}^c$ be the algebra generated by elements $\{\tilde{D}_{0,l} \mid l \geq 1\}$, $\{\tilde{D}_{\pm 1,k} \mid k \geq 0\}$ and $\{\tilde{c}_i \mid i \geq 0\}$ subject to the following set of relations :

$$(3.5) \quad [\tilde{D}_{0,l}, \tilde{D}_{0,k}] = 0, \quad \forall l, k \geq 1,$$

$$(3.6) \quad [\tilde{D}_{0,l}, \tilde{D}_{1,k}] = \tilde{D}_{1,l+k-1}, \quad [\tilde{D}_{-1,k}, \tilde{D}_{0,l}] = \tilde{D}_{-1,l+k-1} \quad \forall l \geq 1, k \geq 0,$$

$$(3.7) \quad (3[\tilde{D}_{1,2}, \tilde{D}_{1,1}] - [\tilde{D}_{1,3}, \tilde{D}_{1,0}] + [\tilde{D}_{1,1}, \tilde{D}_{1,0}]) + \kappa(\kappa - 1)(\tilde{D}_{1,0}^2 + [\tilde{D}_{1,1}, \tilde{D}_{1,0}]) = 0$$

$$(3.8) \quad (3[\tilde{D}_{-1,2}, \tilde{D}_{-1,1}] - [\tilde{D}_{-1,3}, \tilde{D}_{-1,0}] + [\tilde{D}_{-1,1}, \tilde{D}_{-1,0}]) + \kappa(\kappa - 1)(-\tilde{D}_{1,0}^2 + [\tilde{D}_{-1,1}, \tilde{D}_{-1,0}]) = 0$$

$$(3.9) \quad [\tilde{D}_{1,0}, [\tilde{D}_{1,0}, \tilde{D}_{1,1}]] = 0, \quad [\tilde{D}_{-1,0}, [\tilde{D}_{-1,0}, \tilde{D}_{-1,1}]] = 0,$$

$$(3.10) \quad [\tilde{D}_{-1,k}, \tilde{D}_{1,l}] = \tilde{E}_{k+l}, \quad l, k \geq 0,$$

where the $\tilde{E}_l$ are defined by the formula (1.7).

Theorem 3.2. *The assignment $\tilde{D}_{0,l} \mapsto D_{0,l}$, $\tilde{D}_{\pm 1,k} \mapsto D_{\pm 1,k}$ for $l \geq 1, k \geq 0$ and $\tilde{c}_i \mapsto c_i$ for $i \geq 0$ induces an isomorphism of F -algebras*

$$\phi : \widetilde{\mathbf{SH}}^c \xrightarrow{\sim} \mathbf{SH}^c.$$

Coupled with the Theorems in Section 2.3., this provides a potential 'generators and relations' approach to the study of the category of admissible modules over the W -algebras $W_k(\mathfrak{gl}_r)$.

4. PROOF OF THEOREM 3.1

4.1. Let us first observe that ϕ is a well-defined algebra map, i.e. that relations (3.1–3.4) hold in $\mathbf{SH}^+$. For (3.1, 3.2) this follows from the definition of $\mathbf{SH}^+$ and [SV, (1.38)]. Equation (3.3) may be checked directly, e.g. from the Pieri rules (see [SV, (1.26)]), or from the shuffle realization of $\mathbf{SH}^>$ (see 4.2. below). As for equation (3.4), we have by [SV, (1.35)], $[[D_{1,1}, D_{1,0}], D_{1,0}] = [D_{2,0}, D_{1,0}] = 0$. The map ϕ is surjective by construction; in the rest of the proof, we show that it is injective as well.

4.2. Using relation (3.2) it is easy to see that any monomial in the generators $\tilde{D}_{0,l}, \tilde{D}_{1,k}$ may be expressed as a linear combination of similar monomials, in which all $\tilde{D}_{0,l}$ appear on the right of all $\tilde{D}_{1,k}$. Hence the multiplication map $\widetilde{\mathbf{SH}}^> \otimes \widetilde{\mathbf{SH}}^0 \rightarrow \widetilde{\mathbf{SH}}^+$ is surjective. Since ϕ clearly restricts to an isomorphism $\widetilde{\mathbf{SH}}^0 \simeq \mathbf{SH}^0$ we only have to show, by (1.3), that ϕ restricts to an isomorphism $\widetilde{\mathbf{SH}}^> \simeq \mathbf{SH}^>$. Our strategy will be to construct a suitable filtration on $\widetilde{\mathbf{SH}}^>$ mimicking the order filtration of $\mathbf{SH}^>$ and to pass to the associated graded algebras.

4.3. We begin by proving directly, using the shuffle realization of $\mathbf{SH}^>$, that ϕ is an isomorphism in ranks one and two. This is obvious in rank one since ϕ is a graded map and the only relation in rank one is (3.2).

Suppose $\sum \alpha_i D_{1,k_i} D_{1,l_i} = 0$ is a relation in rank two. The shuffle realization then implies $\sum \alpha_i z^{k_i} \star z^{l_i} = 0$ so that

$$h(z_1 - z_2) \left(\sum \alpha_i z_1^{k_i} z_2^{l_i} \right) = h(z_2 - z_1) \left(\sum \alpha_i z_1^{l_i} z_2^{k_i} \right).$$

Therefore $\sum \alpha_i z_1^{k_i} z_2^{l_i} = h(z_2 - z_1)P(z_1, z_2)$ where $P(z_1, z_2)$ is some symmetric polynomial in z_1, z_2 . Hence $\sum \alpha_i z_1^{k_i} z_2^{l_i}$ is a linear combination of polynomials of the form $h(z_2 - z_1)(z_1^k z_2^l + z_1^l z_2^k)$ so that $\sum \alpha_i D_{1,k_i} D_{1,l_i}$ is a linear combination of expressions of the form

$$(4.1) \quad \begin{aligned} & 3[D_{1,l+2}, D_{1,k+1}] - 3[D_{1,l+1}, D_{1,k+2}] - [D_{1,l+3}, D_{1,k}] + [D_{1,l}, D_{1,k+3}] + [D_{1,l+1}, D_{1,k}] - [D_{1,l}, D_{1,k+1}] \\ & + \kappa(\kappa - 1)(D_{1,k} D_{1,l} + D_{1,l} D_{1,k} + [D_{1,l+1}, D_{1,k}] - [D_{1,l}, D_{1,k+1}]). \end{aligned}$$

If I denotes the image of (3.3) under the action of $F[ad\tilde{D}_{0,2}, ad\tilde{D}_{0,3}, \dots]$ then using (3.2) we see that each such expression lies in $\phi(I)$ so that ϕ is indeed an isomorphism in rank two.

We remark that the relations (4.1) may be written in a more standard way using the generating functions $D(z) = \sum_l D_{1,l} z^{-l}$ as follows :

$$(4.2) \quad k(z - w)D(z)D(w) = -k(w - z)D(w)D(z)$$

where $k(u) = (u - 1 + \kappa)(u + 1)(u - \kappa) = -h(-u)$. In particular, the defining relation (3.3) may be replaced by the above (4.2), of which it is a special case.

4.4. We now turn to the definition of the analog, on $\widetilde{\mathbf{SH}}^>$, of the order filtration on $\mathbf{SH}^>$. We will proceed by induction on the rank r . For $r = 1, d \geq 0$, we set

$$\widetilde{\mathbf{SH}}^>[1, \leq d] = \bigoplus_{k \leq d} F\tilde{D}_{1,k}.$$

Assuming that $\widetilde{\mathbf{SH}}^>[r', \leq d']$ has been defined for all $r' < r$ we let $\widetilde{\mathbf{SH}}^>[r, \leq d]$ be the subspace spanned by all products

$$\widetilde{\mathbf{SH}}^>[r', \leq d'] \cdot \widetilde{\mathbf{SH}}^>[r'', \leq d''], \quad r' + r'' = r, d' + d'' = d$$

and by the spaces

$$ad(\tilde{D}_{1,l})(\widetilde{\mathbf{SH}}^>[r - 1, \leq d - l + 1]), \quad l = 0, \dots, d + 1.$$

From the above definition, it is clear that $\widetilde{\mathbf{SH}}^>$ is a $\mathbb{Z}$ -filtered algebra. Note that it is not obvious at the moment that $\widetilde{\mathbf{SH}}^>[r, \leq d] = \{0\}$ for $d < 0$. Because the associated graded $gr\mathbf{SH}^>$ is commutative, it follows by induction on the rank r that $\phi : \widetilde{\mathbf{SH}}^> \rightarrow \mathbf{SH}^>$ is a morphism of filtered algebras. We denote by $gr\widetilde{\mathbf{SH}}^>$ the associated graded of $\widetilde{\mathbf{SH}}^>$ and we let $\bar{\phi} : gr\widetilde{\mathbf{SH}}^> \rightarrow gr\mathbf{SH}^>$ be the induced map. The map $\bar{\phi}$ is graded with respect to both rank and order. Moreover $\bar{\phi}$ is an isomorphism in ranks 1 and 2 (indeed, that the filtration as defined above coincides with the order filtration in rank 2 can be seen directly from [SV, (1.84)]). The rest of the proof of Theorem 3.1 consists in checking that $\bar{\phi}$ is an isomorphism. Once more, we will argue by induction. So in the remainder of the proof, we fix an integer $r \geq 3$ and assume that $\bar{\phi}$ is an isomorphism in ranks $r' < r$.

4.5. By our assumption above, the algebra $gr\widetilde{\mathbf{SH}}^>$ is commutative in ranks less than r , that is $ab = ba$ whenever $rank(a) + rank(b) < r$. Our first task is to extend this property to the rank r .

Lemma 4.1. *The algebra $gr\widetilde{\mathbf{SH}}^>$ is commutative in rank r .*

Proof. We have to show that for $a \in \widetilde{\mathbf{SH}}^>[r_1, \leq d_1], b \in \widetilde{\mathbf{SH}}^>[r_2, \leq d_2]$ and $r_1 + r_2 = r$ we have

$$(4.3) \quad [a, b] \in \widetilde{\mathbf{SH}}^>[r, \leq d_1 + d_2 - 1].$$

We argue by induction on r_1 . If $r_1 = 1$ then (4.3) holds by definition of the filtration. Now let $r_1 > 1$ and let us further assume that (4.3) is valid for all r'_1, r'_2 with $r'_1 + r'_2 = r$ and $r'_1 < r_1$. We will now prove (4.3) for r_1, r_2 , thereby completing the induction step. According to the definition of the filtration, there are two cases to consider :

Case 1) We have $a = a_1 a_2$ with $a_1 \in \widetilde{\mathbf{SH}}^>[s', \leq d']$, $a_2 \in \widetilde{\mathbf{SH}}^>[s'', \leq d'']$ such that $s' + s'' = r_1$, $d' + d'' = d_1$. Then $[a, b] = a_1[a_2, b] + [a_1, b]a_2$. By our induction hypothesis on r , $[a_2, b] \in \widetilde{\mathbf{SH}}^>[s'' + r_2, \leq d'' + d_2 - 1]$ hence $a_1[a_2, b] \in \widetilde{\mathbf{SH}}^>[r, \leq d_1 + d_2 - 1]$. The term $[a_1, b]a_2$ is dealt with in a similar fashion.

Case 2) We have $a = [\tilde{D}_{1,l}, a']$ with $a' \in \widetilde{\mathbf{SH}}^>[r_1 - 1, \leq d_1 - l + 1]$. Then $[a, b] = [[\tilde{D}_{1,l}, a'], b] = [\tilde{D}_{1,l}, [a', b]] - [a', [\tilde{D}_{1,l}, b]]$. By our induction hypothesis on r , $[a', b] \in \widetilde{\mathbf{SH}}^>[r_1 + r_2 - 1, \leq d_1 + d_2 - l]$ hence $[\tilde{D}_{1,l}, [a', b]] \in \widetilde{\mathbf{SH}}^>[r, \leq d_1 + d_2 - 1]$. Similarly, $[\tilde{D}_{1,l}, b] \in \widetilde{\mathbf{SH}}^>[r_2 + 1, \leq d_2 + l - 1]$. The inclusion $[a', [\tilde{D}_{1,l}, b]] \in \widetilde{\mathbf{SH}}^>[r, \leq d_1 + d_2 - 1]$ now follows from the induction hypothesis on r_1 .

We are done. $\square$

4.6. We now focus on the filtered piece of order ≤ 0 of $\widetilde{\mathbf{SH}}^>$. We inductively define elements $\tilde{D}_{l,0}$ for $l \geq 2$ by

$$\tilde{D}_{l,0} = \frac{1}{l-1}[\tilde{D}_{1,1}, \tilde{D}_{l-1,0}].$$

From [SV, (1.35)] we have $\phi(\tilde{D}_{l,0}) = D_{l,0}$. Since we are assuming that $\bar{\phi}$ is an isomorphism in ranks less than r , we have $[\tilde{D}_{l,0}, \tilde{D}_{l',0}] = 0$ whenever $l + l' < r$.

Lemma 4.2. *We have $[\tilde{D}_{l,0}, \tilde{D}_{l',0}] = 0$ for $l + l' = r$.*

Proof. If $r = 3$ this reduces to the cubic relation (3.4). For $r = 4$ we have to consider

$$\begin{aligned} [\tilde{D}_{3,0}, \tilde{D}_{1,0}] &= \frac{1}{2}[[\tilde{D}_{1,1}, \tilde{D}_{2,0}], \tilde{D}_{1,0}] \\ &= \frac{1}{2}[\tilde{D}_{1,1}, [\tilde{D}_{2,0}, \tilde{D}_{1,0}]] - \frac{1}{2}[\tilde{D}_{2,0}, [\tilde{D}_{1,1}, \tilde{D}_{1,0}]] \\ &= -\frac{1}{2}[\tilde{D}_{2,0}, \tilde{D}_{2,0}] = 0. \end{aligned}$$

Now let us fix l, l' with $l + l' = r$. We have

$$\begin{aligned} (\tilde{D}_{l,0}, \tilde{D}_{l',0}) &= \frac{1}{l-1}[[\tilde{D}_{1,1}, \tilde{D}_{l-1,0}], \tilde{D}_{l',0}] \\ (4.4) \quad &= \frac{1}{l-1}[\tilde{D}_{1,1}, [\tilde{D}_{l-1,0}, \tilde{D}_{l',0}]] - \frac{1}{l-1}[\tilde{D}_{l-1,0}, [\tilde{D}_{1,1}, \tilde{D}_{l',0}]] \\ &= -\frac{l'}{l-1}[\tilde{D}_{l-1,0}, \tilde{D}_{l'+1,0}]. \end{aligned}$$

If $r = 2k$ is even then by repeated use of (4.4) we get

$$[\tilde{D}_{l,0}, \tilde{D}_{l',0}] = c[\tilde{D}_k, \tilde{D}_k] = 0$$

for some constant c . Next, suppose that $r = 2k + 1$ is odd, with $k \geq 2$. Applying $ad(\tilde{D}_{1,1})$ to $[\tilde{D}_{k+1,0}, \tilde{D}_{k-1,0}] = 0$ yields the relation

$$(4.5) \quad (k+1)[\tilde{D}_{k+2,0}, \tilde{D}_{k-1,0}] + (k-1)[\tilde{D}_{k+1,0}, \tilde{D}_{k,0}] = 0.$$

Similarly, applying $ad(\tilde{D}_{2,1})$ to $[\tilde{D}_{k,0}, \tilde{D}_{k-1,0}] = 0$ and using the relation $[D_{k,1}, D_{l,0}] = klD_{l+k,0}$ in $\mathbf{SH}^>$ (see [SV, (1.91), (8.47)]) we obtain the relation

$$(4.6) \quad k[\tilde{D}_{k+2,0}, \tilde{D}_{k-1,0}] + (k-1)[\tilde{D}_{k,0}, \tilde{D}_{k+1,0}] = 0.$$

Equations (4.5) and (4.6) imply that $[\tilde{D}_{k+2,0}, \tilde{D}_{k-1,0}] = [\tilde{D}_{k+1,0}, \tilde{D}_{k,0}] = 0$. The general case of $[\tilde{D}_{l,0}, \tilde{D}_{l',0}] = 0$ is now deduced, as in the case $r = 2k$, from repeated use of (4.4). $\square$

Note that Lemma 4.2 above implies that $\widetilde{\mathbf{SH}}^>[r, \leq -1] = \{0\}$.

4.7. Recall that $gr\mathbf{SH}^>$ is a free polynomial algebra in generators in the generators $D'_{s,d}$ for $s \geq 1, d \geq 0$. In order to prove that $\bar{\phi}$ is an isomorphism in rank r , it suffices, in virtue of Lemma 4.1, to show that the factor space

$$U_{r,d} = gr\mathbf{SH}^>[r,d] / \left\{ \sum_{\substack{r'+r''=r \\ d'+d''=d}} gr\mathbf{SH}^>[r',d'] \cdot gr\mathbf{SH}^>[r'',d''] \right\}$$

is one dimensional for any $d \geq 0$. Let us set, for any $s \geq 1, d \geq 0$

$$\tilde{D}'_{s,d} = ad(\tilde{D}_{0,2})^d(\tilde{D}_{s,0}) \in \mathbf{SH}^>[s, \leq d].$$

We will denote by the same symbol $\tilde{D}'_{s,d}$ the corresponding element of $gr\mathbf{SH}^>[s,d]$. Note that $\tilde{D}'_{s,0} = \tilde{D}_{s,0}$. We claim that in fact $U_{r,d} = F\tilde{D}'_{r,d}$. Observe that $\phi(\tilde{D}'_{s,d}) = D'_{s,d}$ for any s, d , hence $\tilde{D}'_{s,d} \in U_{s,d}$ for any $s \leq r, d \geq 0$. Moreover, by our general induction hypothesis on r we have $U_{s,d} = F\tilde{D}'_{s,d}$ for any $s < r$ and $d \geq 0$.

We will prove that $U_{r,d} = F\tilde{D}'_{r,d}$ by induction on d . For $d = 0$, this comes from Lemma 4.2. So fix $d > 0$ and let us assume that $U_{r,l} = F\tilde{D}'_{r,l}$ for all $l < d$. By definition of the filtration on $\mathbf{SH}^>$, $U_{r,d}$ is linearly spanned by the classes of the elements

$$[\tilde{D}_{1,0}, \tilde{D}'_{r-1,d+1}], [\tilde{D}_{1,1}, \tilde{D}'_{r-1,d}], \dots, [\tilde{D}_{1,d+1}, \tilde{D}'_{r-1,0}].$$

By our induction hypothesis on d , the elements

$$[\tilde{D}_{1,0}, \tilde{D}'_{r-1,d}], [\tilde{D}_{1,1}, \tilde{D}'_{r-1,d-1}], \dots, [\tilde{D}_{1,d}, \tilde{D}'_{r-1,0}]$$

all belong to $F\tilde{D}'_{r,d-1} \oplus \mathbf{SH}^>[r, \leq d-2]$. Applying $ad(\tilde{D}_{0,2})$, we see that

$$(4.7) \quad [\tilde{D}_{1,0}, \tilde{D}'_{r-1,d+1}] + [\tilde{D}_{1,1}, \tilde{D}'_{r-1,d}], \dots, [\tilde{D}_{1,d}, \tilde{D}'_{r-1,1}] + [\tilde{D}_{1,d+1}, \tilde{D}'_{r-1,0}]$$

all belong to $F\tilde{D}'_{r,d} \oplus \mathbf{SH}^>[r, \leq d-1]$. Next, applying $ad(\tilde{D}_{0,d+2})$ to the equality $[\tilde{D}_{1,0}, \tilde{D}_{r-1,0}] = 0$ yields

$$[\tilde{D}_{1,0}, \tilde{D}_{r-1,d+1}] + [\tilde{D}_{1,d+1}, \tilde{D}_{r-1,0}] = 0$$

which implies, by (1.4), that

$$(4.8) \quad [\tilde{D}_{1,0}, \tilde{D}'_{r-1,d+1}] + r^d[\tilde{D}_{1,d+1}, \tilde{D}_{r-1,0}] \in [\tilde{D}_{1,0}, \mathbf{SH}^>[r-1, \leq d]] \subseteq \mathbf{SH}^>[r, \leq d-1].$$

The collection of inclusions (4.7), (4.8) may be considered as a system of linear equations in $U_{r,d}$ modulo $F\tilde{D}'_{r,d}$ in the variables $[\tilde{D}_{1,0}, \tilde{D}'_{r-1,d+1}], \dots, [\tilde{D}_{1,d+1}, \tilde{D}'_{r-1,0}]$ whose associated matrix is

$$M = \begin{pmatrix} 1 & 0 & \cdots & 0 & 1 \\ 1 & 1 & \cdots & 0 & 0 \\ 0 & 1 & \ddots & \vdots & \vdots \\ \vdots & \vdots & \ddots & 1 & 0 \\ 0 & 0 & \cdots & 1 & -r^d \end{pmatrix}$$

is invertible. We deduce that $[\tilde{D}_{1,0}, \tilde{D}'_{r-1,d+1}], \dots, [\tilde{D}_{1,d+1}, \tilde{D}'_{r-1,0}]$ all belong to the space $F\tilde{D}'_{r,d} \oplus \mathbf{SH}^>[r, \geq d-1]$ as wanted. This closes the induction step on d . We have therefore proved that $U_{r,d} = F\tilde{D}'_{r,d}$ for all $d \geq 0$, and hence that $\bar{\phi}$ and ϕ is an isomorphism in rank r . This closes the induction step on r . Theorem 3.1 is proved. $\square$

Acknowledgements

We would like to thank P. Etingof for helpful discussions. The first author would like to thank the mathematics department of Orsay University for its hospitality and the MIT-France program for supporting his stay.

REFERENCES

- [A] T. Arakawa, *Representation theory of W -algebras*, Invent. Math. **169** (2007), 219–320.
- [C] I. Cherednik, *Double affine Hecke algebras*, London Mathematical Society Lecture Notes Series, **319**, Cambridge University Press, Cambridge (2005).
- [FO] B. Feigin, A. Odesskii, *Vector bundles on an elliptic curve and Sklyanin*, Topics in quantum groups and finite type invariants, 65–84, Amer. Math. Soc. Transl. Ser. 2, **185**, (1998).
- [FKRW] E. Frenkel, V. Kac, A. Radul, W. Weiqiang, *$W_{1+\infty}$ and $W(\mathfrak{gl}_N)$ with central charge N* , Commun. Math. Phys., **170** (1995), 337–357.
- [MO] D. Maulik, A. Okounkov, *private communication*.
- [SV] O. Schiffmann, E. Vasserot, *Cherednik algebras, W -algebras and the equivariant cohomology of the moduli space of instantons on $\mathbb{A}^2$* , preprint arXiv:1202.2756 (2012)
- [S] R. Stanley, *Some combinatorial properties of Jack symmetric functions*, Advances in Math., **77**, 76–115 (1989).

N. Arbesfeld, nma@mit.edu,
 Massachusetts Institute of Technology Cambridge, MA 02139, USA.

O. Schiffmann, olivier.schiffmann@math.u-psud.fr,
 Département de Mathématiques, Université de Paris-Sud, Bâtiment 425 91405 Orsay Cedex, FRANCE.